

BASF Palladium Catalyst Kit



METALS • INORGANICS • ORGANOMETALLICS • CATALYSTS • LIGANDS • NANOMATERIALS • CUSTOM SYNTHESIS • CGMP FACILITIES		
96-6719 HAZ	BASF Palladium Catalyst Kit Product offered is commercial grade, sold in collaboration with BASF for research purposes only. Components also available for individual sale. Contains the smallest unit size of each of the following item	ns:
46-1707	Palladium, 20% on activated carbon (Pearlman's catalyst), unreduced, 50% water wet paste (Escat™ 19 (7440-05-3) black pwdr. (d50=24 µm); SA: 850m²/g	5 1) 5g 25g
Technical Note: 1. Escat™ 1951 catalyst is recommended for a broad range of reactions common to palladium on carbon catalysts. Specifically, it is well suited for removal of protecting groups such as benzyl, FMOC and others.		
46-1710	Palladium, 0.6% on activated carbon, 50% water-wet paste (NanoSelect LF 100) $(7440$ -05-3) black solid (d50=25 μ m)	5g 25g
 Technical Note: 1. NanoSelect LF 100 is a lead-free, water-wet, catalyst containing metal crystallites sizes of around 7 nm, and a mean particle size of 25 microns. The metal crystallites are supported on a carbon powder. The presence of nanometer-sized metal particles greatly increases the metal surface area available per gram of catalyst, and boosts catalytic activity. The catalyst is recommended for use in hydrogenation reactions leading to the partial reduction of functional groups. It is specifically suited for the selective hydrogenation of alkynes to alkenes, with a high selectivity for cis-alkenes. 		
46-1901	Palladium, 5% on activated peat carbon, reduced, 50% water wet paste (Escat™ 1621) (7440-05-3) black pwdr. (d50=15 μm); SA: 850m²/g	10g 50g
Technical Note: 1. Escat™ 1621 catalyst is recommended for a broad range of reactions common to palladium on carbon catalysts, such as hydrogenolysis under hydrogen transfer conditions.		
46-1902 HAZ	Palladium, 5% on activated wood carbon, reduced, dry (Escat [™] 1431) (7440-05-3) black pwdr. (d50=18 µm); SA: 900m²/g	10g 50g
Technical Note: 1. Escat™ 1431 catalyst is recommended for a broad range of reactions common to palladium on carbon catalysts, where water is detrimental to the selectivity of the reaction. Active over a wide range of conditions.		
46-1903	Palladium, 5% on activated wood carbon, reduced, 50% water wet paste (Escat™ 1421) (7440-05-3) black pwdr. (d50=18 μm); SA: 900m²/g	10g 50g
Technical Note: 1. Escat™ 1421 catalyst is recommended for a broad range of reactions commonly catalyzed by palladium on carbon.		
46-1905	Palladium, 10% on activated wood carbon, reduced, 50% water wet (Escat™ 1931) (7440-05-3) black pwdr. (d50=38 μm); SA: 1500m²/g	10g 50g
	te: [™] 1931 catalyst is recommended for a broad range of reactions commonly catalyzed by palladium on carbon. S for removal of protecting groups such as benzyl, FMOC and others.	Specifically, it is well
46-1904	Palladium, 5% on activated wood carbon, unreduced, 50% water wet paste (Escat™ 1471) (7440-05-3) black pwdr. (d50=18 µm); SA: 900m²/g	10g 50g
Technical Note: 1. Escat™ 1471 catalyst is recommended for a broad range of reactions common to palladium on carbon catalysts. Specifically, it is well suited for hydrogenolysis reactions using molecular hydrogen. Active over a wide range of conditions.		
46-1906	Palladium, 10% on activated wood carbon, unreduced, 50% water wet (Escat™ 1921) (7440-05-3) black pwdr. (d50=38 μm); SA: 1500m²/g	10g 50g
Technical Note: 1. Escat™ 1921 catalyst is recommended for a broad range of reactions common to palladium on carbon catalysts. Specifically, it is well suited for removal of protecting groups such as benzyl, FMOC and others.		
46-1951	Palladium, 5% on alumina powder, reduced, dry (Escat™ 1241) (7440-05-3) gray pwdr. (d50=70 μm); SA: 110m²/g	5g 25g

Technical Note:

1. Escat™ 1241 catalyst is recommended for selective hydrogenation reactions such as alkyne reduction in the presence of carboxylic groups. The particle size of the catalyst is ideal for allowing fast separation from the reaction mixture.

Visit www.strem.com for new product information and a searchable catalog.

Strem Chemicals, Inc. 7 Mulliken Way Newburyport, MA 01950 U.S.A Tel: 978.499.1600 Fax: 978.465.3104 Email: info@strem.com

Strem Chemicals, Inc. 15, rue de l'Atome Zone Industrielle 67800 BISCHHEIM France Tel: (33) 03 88 62 52 60 Fax: (33) 03 88 62 26 81 Email: info.europe@strem.com Strem Chemicals, Inc. Postfach 1215 77672 KEHL Germany Tel: 0 78 51/ 7 58 79

Email: info.europe@strem.com

Strem Chemicals UK Ltd.

An Independent Distributor of Strem Chemicals Products Newton Hall, Town Street Newton, Cambridge England CB22 7ZE Tel: +44 (0)1223 873 028

Fax: +44 (0)1223 870207 Email: enquiries@strem.co.uk